

## Additional References

- Bradley, M. C., Leidich, S., Isham, N., Elewski, B. E., and Ghannoum, M. 1999. Antifungal susceptibilities and genetic relatedness of serial *Trichophyton rubrum* isolates from patients with onychomycosis of the toenail. *Mycoses.* 42 (Supp. 2):105-110.
- Ghannoum, M. A., Espinel-Ingroff, A., Pfaller, M. A., et al. An Intra- and Inter-Laboratory Study of an Antifungal susceptibility Method for the Testing of Dermatophytes.
- Jessup, C. J., Warner, J., Isham, N., Hasan, I., and Ghannoum, M. 2000. Antifungal Susceptibility Testing of Dermatophytes: Establishing a Medium for Inducing Conidial Growth and Evaluation of Susceptibility of Clinical Isolates. *J. Clin. Microbiol.* 38(1):341-344.
- Martin, A. G. and Kobayashi, G. S. "Superficial fungal infections: dermatophytosis, tinea nigra, piedra" in Freedberg, I. M., et al. (ed) (1999), *Fitzpatrick's Dermatology in General Medicine*, New York: McGraw-Hill, Chapter 206, 2349-2351.
- Mukherjee, P., Leidich, S., and Isham, N., et al. 2003. Clinical *Trichophyton rubrum* Strain Exhibiting Primary Resistance to Terbinafine. *Antimicrob. Agents Chemother.* 47:82-86.
- Norris, H., Elewski, B., and Ghannoum, M. 1999. Optimal growth conditions for the determination of antifungal susceptibility of three species of dermatophytes with the use of a microdilution method. *J. Am. Acad. Dermatol.* 40(6)(2):S9-S-13.
- Roberts, D. T. and Bilsland, D. J. "Tinea pedis and skin dermatophytosis" in Lebwohl, M. et al. (ed) (2002), *Treatment of skin disease. Comprehensive therapeutic strategies*, Philadelphia: Mosby, Section 200, 624-626.